



# Nanoparticle-Enriched Edible Coatings with Plant Extracts and Essential Oils for Extending Postharvest Quality of Fruits and Vegetables

Praveen Gidagiri <sup>a\*</sup>, Lekshmi P. R. G. <sup>a</sup> and Shameena S. <sup>a</sup>

<sup>a</sup> Department of Postharvest Management, College of Agriculture, Vellayani, Kerala Agricultural University, 695522, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2025/v28i11881>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129675>

Review Article

Received: 07/11/2024  
Accepted: 10/01/2025  
Published: 13/01/2025

## ABSTRACT

Fruits and vegetables are affected by postharvest diseases and physiological processes such as transpiration and respiration after harvest, leading to significant quality losses during transportation and storage. To control postharvest diseases and extend shelf life, synthetic fungicides are commonly used. However, the extensive use of these chemicals raises concerns about potential health risks, environmental impact, and the development of fungicide-resistant pathogens. Therefore, alternative approaches, such as edible coatings with plant extracts, essential oils and nanoparticles loaded with plant extracts are explored to address these challenges while maintaining quality and safety of fresh fruits and vegetables. Nanoparticles loaded with essential oils or plant extracts are synthesized using ionic gelation method, spontaneous emulsification,

\*Corresponding author: E-mail: [praveen-2021-22-007@student.kau.in](mailto:praveen-2021-22-007@student.kau.in);

**Cite as:** Gidagiri, Praveen, Lekshmi P. R. G., and Shameena S. 2025. "Nanoparticle-Enriched Edible Coatings With Plant Extracts and Essential Oils for Extending Postharvest Quality of Fruits and Vegetables". *Journal of Advances in Biology & Biotechnology* 28 (1):290-300. <https://doi.org/10.9734/jabb/2025/v28i11881>.

phase inversion temperature, emulsion inversion point, and solvent displacement methods etc. Nanoparticle enriched edible coating delivers plant compound gradually over time. This controlled release provides a consistent and long-lasting effect, effectively inhibiting microbial growth, slowing nutrient degradation, and reducing oxidative stress on the produce. The nanoscale size of these particles increases their surface area, enhancing their adhesion to the surface of fruits or vegetables. This results in a more efficient barrier against moisture loss and gas exchange, further preserving the quality and extending the shelf life of the produce. The synergistic effects of the nano-reinforced matrices show promising results in reducing post-harvest losses while maintaining food safety standards. This innovative approach holds immense potential for revolutionizing the postharvest management of fruits and vegetables by offering a safer, more sustainable alternative to synthetic fungicides. Future research should focus on optimizing the formulation and scalability of nanoparticle-enriched edible coatings, evaluating their efficacy across diverse produce types, and assessing their long-term safety and environmental impact. This review examines the current developments in edible coating technology, focusing on various plant extracts, essential oils, and nanoparticles loaded with plant extracts enriched edible coating for extending shelf life and preserving the quality of fruits and vegetables.

*Keywords: Nanoparticle-enhanced coatings; plant extracts; food preservation; postharvest technology.*

## 1. INTRODUCTION

India is the second largest producer of fruits and vegetables and plays a major role in food, nutritional and economic security of the country. The area under fruit cultivation is 69.3 lakh ha with a production of 102.481 million tons and vegetables production is 200.445 million tons from an area of 102.60 lakh ha (NHB, 2020). "However, major problem in fruits and vegetables is their higher perishability leading to postharvest losses and short shelf life. The postharvest loss in India is reported to be around 8-18% (APEDA, 2019) because of high moisture content (75–95%) in fruits and vegetables, leading to dehydration, mechanical injury, environmental stress, pathological breakdown, poor postharvest management, inadequate cold storage, processing facilities etc" (Otoni et al., 2017).

"Preserving postharvest quality is a critical concern for fruits and vegetables growers to ensure efficient handling, transportation, and marketing of fresh produce" (Dave et al., 2017). Various chemical compounds are commonly used to control postharvest diseases and to extend the shelf life of fruits and vegetables, some of these substances pose risks to human health. "Additionally, the widespread use of fungicides and synthetic chemicals has contributed to the emergence of resistant strains of pathogens, accompanied by the accumulation of harmful residues" (Khaliq et al., 2019). "With growing consumer demand for safe, high-quality produce, minimizing postharvest losses has become increasingly important" (Adekunle et al.,

2021). "To address these challenges, the application of edible coatings has been explored as a sustainable solution. These eco-friendly, thin-layer substances are applied to fresh fruits and vegetables to reduce postharvest losses and maintain their quality" (Dhall et al., 2013).

Edible coatings can serve as carriers for active ingredients that help to extend the postharvest life of fruits and vegetables, reduce the risk of pathogen infestation on food surfaces, and slow down other undesirable changes. This can be accomplished using different application techniques such as dipping, spraying, or brushing. The coating formulation may consist of a single ingredient or a blend of various components, including polysaccharides (e.g., starch, cellulose derivatives), proteins (e.g., gelatin, whey protein), lipids (e.g., waxes, fatty acids), and additional additives like plasticizers, antimicrobials (Sharma et al., 2024). "Polysaccharide-based edible coatings are widely used to prolong the storage life of fruits" (Cazon et al., 2017; Liu et al., 2021). "These coatings provide protection by minimizing moisture loss, regulating respiration and gas exchange, and inhibiting the activity of reactive oxygen species" (Aloui, et al., 2016). Enhancing these coatings with plant extracts or essential oils increases their effectiveness while making them more environmentally friendly. Additionally, these coatings offer several benefits, such as being cost-effective and readily available. "The effectiveness of plant-based edible coatings or those enriched with plant extracts largely depends on the active compounds in the

extracts, which are generally recognized as safe for human consumption and possess strong antioxidant and antimicrobial properties” (Flores-Lopez et al., 2016).

“The incorporation of nanoparticles into edible coatings enhances their functional, nutritional, organoleptic, and mechanical properties, presenting a novel approach to maintaining product quality. Research has demonstrated that adding nanoparticles to edible coatings improves their effectiveness in preserving the quality of fresh produce, provides antibacterial properties, and enhances food safety during storage” (Martelli et al., 2013). “Nanoparticles are also employed in food science to improve the quality and safety of various products, including shrimp” (Baek et al., 2021), seafood (Ozuna-Valencia et al., 2024), and minimally processed pomegranate (Anean et al., 2023). The integration of plant-based materials into edible coatings represents a recent and innovative technology. These coatings, being entirely organic, not only preserve the postharvest quality of fruits and vegetables but also ensure consumer safety. This review highlights the potential of edible coatings enriched with plant extracts, essential oils, and nanoparticles loaded with these bioactive compounds. It examines their impact on various physical, biochemical, antioxidant, and enzymatic parameters during the postharvest period of fruits and vegetables.

## 2. PLANT EXTRACTS AND ESSENTIAL OILS FOR EDIBLE COATING

“Edible coating formulations with incorporation of plant extracts and essential oils have become increasingly popular nowadays due to its antimicrobial properties that helps to controlling postharvest diseases and slowing the deterioration of fruits and vegetables” (Saxena et al., 2020). “In recent years, there has been growing interest in enhancing the performance of edible coatings for fresh and fresh-cut fruits and vegetables, especially when the coatings fail to achieve all the desired effects. A notable trend involves the incorporation of essential oils and plant extracts into these coatings. Essential oils are naturally occurring volatile compounds with complex structures and strong aromas, produced by aromatic plants as secondary metabolites. These compounds play a crucial role in the plant’s defense mechanisms” (Soleimani, et al., 2023). “Essential oils are widely recognized for their antibacterial, antifungal, and insecticidal properties. Classified as GRAS (Generally

Recognized as Safe) substances, essential oils derived from plant materials are effective against foodborne pathogens and spoilage-causing bacteria” (Hager et al., 2019). Among the essential oils, commonly extracted from spices and incorporated into edible coatings are those from the Myrtaceae family (e.g., clove and tea tree) and the Lauraceae family (e.g., cinnamon). The essential oils used in edible coatings from herbs and grasses are typically obtained from the Lamiaceae family (e.g., thyme, peppermint, rosemary, oregano, and savory), the Asteraceae family (e.g., tarragon), and the Poaceae family (e.g., lemongrass). Additionally, citrus-based essential oils have been applied in edible coatings for fresh and fresh-cut fruits and vegetables, with notable examples coming from the Rutaceae family, such as lemon, lime, and bergamot (Yousuf et al., 2021).

“Plant extracts are complex blends of compounds belonging to various chemical classes that possess bioactive properties. Traditionally, these extracts are obtained from the entire plant or specific parts such as the rhizome, root, stem, leaf, flower, or tissues and cell cultures” (Hao et al., 2015). “The bioactive compounds within these mixtures are extracted from plants or plant materials using different solvents” (Sasidharan et al., 2011). These bioactive compounds, along with other natural substances like metabolites, volatile compounds, and active enzymes produced by selected microorganisms, serve as valuable sources of antifungal agents, making them promising candidates for fungicides to manage post-harvest fungal diseases (Bhutia, 2015). As noted by Gurjar et al. (2012), plant tissues have the capability to produce aromatic secondary metabolites, including phenols, phenolic acids, quinones, flavones, flavonoids, flavonols, tannins, and coumarins.

Moringa extract is a potent source of vitamins, minerals, proteins,  $\beta$ -carotene, phenolic acids, flavonoids, and various other bioactive compounds (Saucedo-Pompa et al., 2018). Its high levels of phenolic acids and flavonoids are largely responsible for its strong antioxidant activity, demonstrated in both in vitro and in vivo studies, as reported by Vongsak et al. (2013). Research has shown that moringa extract exhibits several beneficial properties, including anti-inflammatory, antifungal (Chung et al., 2007), pro-coagulant (Nkurunziza et al., 2009), flocculating (Beltran-Heredia et al., 2009), and antibacterial activities (Doughari et al., 2007).

“Papaya leaves and seeds are rich in proteolytic enzymes (papain and chymopapain), alkaloids (carpain and carpasemine), sulfur-containing compounds (benzyl isothiocyanate), flavonoids, triterpenes, and organic acids. Extracts derived from various papaya tissues have demonstrated bioactive properties. Aqueous extracts from papaya leaves and seeds exhibit antifungal activity against *Colletotrichum gloeosporioides*” (Bautista-Banos et al., 2002). “Additionally, alcoholic extracts from the epicarp, endocarp, roots, and seeds of ripe and unripe papaya fruits possess antidiarrheal, antidysenteric, and antibacterial properties” (Doughari et al., 2007).

The application of moringa leaf extract (2%) and a combination of moringa leaf extract with 1% carboxymethyl cellulose on ‘Fuerte’ and ‘Hass’ avocado fruits significantly reduced respiration rate, moisture loss, and firmness loss during storage, thereby extending their postharvest shelf life (Tesfaya and Magwazaa, 2017). Similarly, postharvest treatments using 10% moringa leaf extract combined with 10% gum arabic or 1% carboxymethyl cellulose effectively controlled *Colletotrichum gloeosporioides* by up to 33%, minimized weight and firmness loss, and delayed colour changes in ‘Maluma’ avocados (Kubheka et al., 2020). The edible coating of guava (cv. Maamoura) with 10% gum arabic and 10% moringa leaf extract effectively reduced postharvest weight loss, decay, and *Rhizopus* rot infection, while delaying fruit softening under cold storage conditions and improving marketability (El-Gioushy et al., 2022). Similarly, the application of 6% moringa leaf extract on strawberry fruits (cv. Chandler) resulted in the lowest respiration rate, ethylene production, malondialdehyde levels, and total soluble solid accumulation compared to untreated fruits during five days of storage at ambient conditions (Shafique et al., 2023).

Tabassum et al. (2018) studied the effects of guava leaf and lemon extracts on the postharvest quality of Sabri bananas under ambient conditions. A combination of 20% guava leaf extract and 15% lemon extract was most effective in maintaining higher firmness, reducing total soluble solids and microbial infection, and extending shelf life to 8.75 days. Meanwhile, 40% guava leaf extract with 5% lemon extract preserved the highest vitamin C (1.83 mg/100g), titratable acidity (0.11%), and reducing sugar (7.13%) after two weeks of storage. Khaliq et al. (2019) examined the effects of aloe vera gel enriched with *Fagonia indica* extract on the

quality and antioxidant activity of sapodilla fruit during postharvest storage at 20 °C for 12 days. Aloe vera gel (50% or 100%) combined with 1% *Fagonia indica* extract significantly reduced weight loss, decay, and soluble solids concentration while preserving firmness, titratable acidity, ascorbic acid, total flavonoids, phenolics, and antioxidant activity. Sensory analysis showed no negative impact on fruit attributes, highlighting the potential of this treatment to extend sapodilla fruit shelf life and maintain quality. Moringa leaf extract combined with carboxymethyl cellulose serves as an eco-friendly edible coating to reduce postharvest losses in “Hass” avocados. A coating of 16 % moringa leaf extract with 5% carboxymethyl cellulose effectively preserved phenolics, flavonoids, and antioxidant activity and firmness during storage at 5.5°C and 90% RH (Ngubane et al., 2024).

### 3. EFFECT OF PLANT EXTRACTS/ESSENTIAL OILS LOADED NANOPARTICLES ON QUALITY AND SHELF LIFE OF FRUITS AND VEGETABLES

Nanotechnology has emerged as a powerful tool for developing edible coatings based on emulsions with enhanced properties and functionalities. It represents a novel approach to improving postharvest management of fruits and vegetables. Incorporating nanoparticles into edible coatings significantly extends the shelf life of fruits compared to coatings made solely from pure polymers (Gad and Zog, 2017). The inclusion of nanoparticles enhances the mechanical and barrier properties of the coatings while also increasing their thermal stability (Shankar and Rhim, 2015).


Most successful applications of nanotechnology in the field of packaging concerns the development of “nanocomposites” (Unalan et al., 2014). In polymer science, while the term “composites” generally refers to mixtures of polymers with inorganic or organic additives having micron-length scale and certain geometries (fibers, flakes, spheres, and particulates), the use of nano-length-scale entities can be more specifically referred to as “nanocomposites.” A new subclass of nanocomposite materials has recently stemmed from the increasing endeavour to replace oil-based polymers with polymers partially or totally obtained from renewable resources. The term

“bionanocomposites” refers to those materials in which the polymer matrix carrying the nanosized fillers is a biopolymer (e.g., polysaccharides, lipids and proteins). Composite materials exhibit stronger physical, chemical, and mechanical properties than their constituent materials. However, conventional composite materials mechanically differ from nanocomposites because of their exceptionally high surface-to-volume ratio and are excellent green technology materials with good biodegradability, biocompatible properties, and the capability to mimic biomaterials.

Banana fruits coated with Cellulose nanofiber based emulsion (0.3%) delayed the ethylene biosynthesis pathway, reduced ethylene and CO<sub>2</sub> production, minimized chlorophyll degradation, weight loss, and firmness of bananas during 10 days of ambient storage (Deng et al., 2017). Banana (Cavendish) fruits coated with chitosan nanoparticles 0.2% has a slower skin discoloration by 2-3 days compared to control treatment under 22 ± 1°C (Esyanti et al., 2019). Jianmei et al. (2019) studied the improvement of banana postharvest quality using a novel soybean protein isolate (SPI) / cinnamaldehyde (CIN) / zinc oxide (ZnO) bionanocomposite coating strategy. Results showed that SPI/CIN/ZnONP nanocomposite coating could effectively delay ripening rate, lowest weight loss (7.83 %), TSS (10.31 %) and high fruit firmness (12.13 N), titratable acidity (65.93 mg malic acid/100 g), sensory quality compared with other SPI-based coating

treatments during the whole storage period (7 days).

Barrera-Necha et al. (2018) studied the synthesis and characterization of chitosan nanoparticles loaded botanical extracts with antifungal activity on *Colletotrichum gloeosporioides*. In this study, chitosan nanoparticles (CSNPs) and chitosan nanoparticles botanical extracts: EMN-CSNPs (extract methanol of nanche added chitosan nanoparticles) were characterized and evaluated *in vitro* on growth of *Colletotrichum gloeosporioides* isolated from Papaya and Soursop. For *in vitro* evaluation, incorporation of EMN to CSNPs improved the control of *C. gloeosporioides* with amycelial growth inhibition of 79% from papaya and 82% from soursop. Odetayo et al. (2022) studied the effect of nanoparticle-enriched coatings on the shelf life of Cavendish bananas. Banana fruits were dipped into: *Aloe vera* 50% (AV), *Moringa oleifera* 10% (MO), *Aloe vera* 50% + chitosan nanoparticle 2% (AV+CN), and *Moringa oleifera* 10% + chitosan nanoparticle 2% (MO+CN). The results showed that adding chitosan nanoparticles to edible coating had a significant impact on the weight loss, firmness, ethylene rate and disease incidence during storage. *Moringa oleifera* + chitosan nanoparticle coating reduced weight loss (23%), ethylene generation (144 µL/kg/h) and disease incidence (58 %) to a larger extent followed by other treatments. In addition, the MO+CN treatment preserved fruit firmness (46 N) and increased the shelf life to 30 days compared to others.



**List 1. Types of base materials used for edible coating**  
(Saxena et al., 2020)

**Table 1. Studies on bionanocomposite edible coatings on different fruits and vegetables**

Sl. No.	Fruit/vegetable	Nanoparticle component	Bioactive compound	Reference
1.	Apple	Zein nanofibers	Curcumin	
<b>Result</b>		The surface was inoculated with <i>Botrytis cineraria</i> and <i>Penicillium expansum</i> ; then, apples were coated by electro-spinning with zein nanotubes and stored for 15 days, revealing the inhibition of microbial growth and increase in the shelf life of apples.		Aytac et al., 2017
2.	Cucumber	Chitosan	Carvacrol	
<b>Result</b>		The combination of pulsed light (12 J/cm <sup>2</sup> ) with the edible coating (0.08% carvacrol) resulted in a strong synergistic effect, with <i>E. coli</i> reduction reaching >5 log cycles.		Tastan et al., 2017
3.	Grape berry	Chitosan	Lemongrass oil	
<b>Result</b>		The use of the nanoemulsion effectively reduced the initial growth of <i>Salmonella typhimurium</i> , total aerobic mesophiles, yeasts and moulds, and showed retention of antioxidant capacity.		Cordoba and Sobral, 2017
4.	Green beans	Chitosan	Mandarin essential Oil	
<b>Result</b>		The combination of the bioactive coating and UV-C treatment reduced the <i>Listeria innocua</i> population and maintained the microbial load at a constant level during storage.		Severino et al., 2014
5.	Fresh-cut apple	Sodium alginate	Lemongrass essential oil	
<b>Result</b>		Nanoemulsion based edible coatings presented higher <i>E. coli</i> inactivation and slower psychrophilic bacteria growth compared to conventional emulsions at the same concentration.		Salvia-Trujillo et al., 2015

“The curcumin-loaded chitosan nanoparticles effectively controlled the anthracnose disease on Papaya under both *in vitro* and *in vivo* assay. The Curcumin alone inhibited the growth of *C. gloeosporioides* at both *in vitro* and *in vivo* assays. However, the Cur-ChNP with formula ratio of 5:5 (w/w), had similar effect with Curcumin treatment to suppress anthracnose disease severity under *in vivo* assay, with the relative disease inhibition rate of 71.19%. This study, demonstrated that Cur-ChNP may be utilized as a potential biocontrol agent against anthracnose disease on papaya” (Suryadi et al., 2019). Vieira et al. (2020) reported that papaya fruits coated with Hydroxypropyl methylcellulose (4%) + silver nanoparticles (0.25%) preserved the quality of papaya evaluated by colour, firmness, lower weight loss, and total soluble solids. Thus, it can be an alternative against fungus development and to increase the shelf life of *papaya cv. Golden*.

Resende et al. (2018) observed that 1% chitosan + 5% CNF (cellulose nanofibril) coating

performed better in reducing fruit mass, firmness loss and maintaining overall quality of strawberry under cold storage (1 ± 1°C). Melo et al. (2020) studied on *In vivo* and *in vitro* antifungal effect of fungal chitosan nanocomposite edible coating against strawberry phytopathogenic fungi. The results obtained in this study demonstrated that the application of chitosan as a gel, or nanocomposite can control the growth and the viability of the strawberry phytopathogenic fungi (*Botrytis cinerea*; *Rhizopus stolonifer* and *Aspergillus niger*). The application of the nanocomposite edible coating *in vitro* and on strawberries artificially contaminated with postharvest fungi, decreased the infection severity caused by these fungi during fruit storage. Therefore, chitosan nanocomposite edible coatings can be a promising strategy to extend the shelf life of strawberries. Nguyen et al. (2020) studied on combination effects of calcium chloride and nano-chitosan on the postharvest quality of strawberry (*Fragaria x ananassa* Duch.). The fruit were dipped in different concentrations of calcium chloride

(1 %, 2 %, 3 %, 4 %) before being coated with 0.2 % nano-chitosan. Among six examined treatments, a combination of 3 % CaCl<sub>2</sub> and nano-chitosan (NCTS) 0.2 % was the most effective one as maintaining the highest score of overall quality index of strawberry stored at 4 °C up to 15 days.

“A bionanocomposite edible coating containing carboxymethyl cellulose (CMC) and cardamom essential oil (CEO) significantly reduced the loss of fruit weight ( $7.32 \pm 2.4\%$ ), firmness (1.3 times), total soluble solids, and titratable acidity of tomatoes by reducing oxidative stress and increasing antioxidant enzymes during 15 days of storage” (Das et al., 2022). Algarni et al. (2022) “studied on effect of chitosan nanoparticles as edible coating on the storability and quality of apricot fruits. During the storage of the apricots, those treated with CHNPs (1%) showed an obvious decrease in decay percent and lipid peroxidation, whereas ascorbic acid and carotenoid content were higher than those in the fruits treated with CH and the untreated fruits (control). The findings of the sensory evaluation revealed a significant difference in the overall acceptability scores and extended their shelf-life up to 9 days at room temperature storage and for 30 days in cold storage between the samples treated with CHNPs and the other samples.”

Mohammadi et al. (2015) “studied on chitosan nanoparticles loaded with *Cinnamomum zeylanicum* essential oil enhance the shelf life of cucumber during cold storage.” Chitosan nanoparticles loaded with CEO significantly decreased both disease severity and incidence of infected cucumbers by *P. drechsleri*, during 7 days of storage at 4 °C followed by 1–2 more days at 20 °C. Furthermore, the CEO-CSNs was more effective than pure CSNs coating at reducing respiration rates, improving the microbiological quality, and preserving the fruit weight and quality of cucumbers during the storage period. These results demonstrate the potential of chitosan nanoparticles containing cinnamon oil extended the postharvest shelf life of cucumber. Emamifar and Bavaisi (2020) reported that strawberries coated with 1.5% sodium alginate + 1.25 g/L nano-ZnO showed the highest antioxidant activity, antimicrobial properties, sensory attributes, and lowest peroxidase activity and extended the storage life of fresh fruits by up to 20 days.

Saekow et al. (2019) “studied on effect of carboxymethyl cellulose coating containing ZnO-

nanoparticles for prolonging the shelf life of persimmon and tomato fruit. The results showed that the application of coating with carboxymethyl cellulose and carboxymethyl cellulose + ZnO-nanoparticles in persimmon and tomato reduced weight loss and respiration rate, and increased fruit firmness when compared with control during storage time. The CMC + ZnONPs coating effectively delayed the disease severity in the inoculated persimmon and tomato”. Maringgal et al. (2021) “studied on effect of kelulut honey nanoparticles (KHNPs) coating on the changes in respiration rate, ascorbic acid, and total phenolic content of papaya (*Carica papaya* L.) during cold storage. Papayas coated with 15 % KHNPs significantly changed during storage, with an increase in CO<sub>2</sub> and a decrease in O<sub>2</sub> and C<sub>2</sub>H<sub>4</sub>, while the ascorbic acid and total phenolic content were maintained. The changes in respiration rate were rather slower for coated papayas when compared to the control.”

#### 4. CONCLUSION

Edible coating is one of the promising technologies for extending the shelf life of fresh fruits and vegetables that help to reduce postharvest losses. Incorporating plant extracts and essential oils into edible coatings has proven effective in extending the postharvest life of various fruits and vegetables. Edible coatings enriched with plant extracts are gaining popularity due to their antioxidant properties, affordability, safety, and easy availability. Recent advancements in postharvest technology have provided new methods for integrating these plant extracts into the edible coating matrix. The inclusion of nanoparticles in edible coatings significantly extends shelf life. Recent studies show that nanoparticles loaded with plant extracts and essential oils play a crucial role in developing next-generation edible coatings with enhanced properties for preserving fresh fruits and vegetables. This emerging technology improves the physical stability and effectiveness of active ingredients within edible coatings, offering the potential to enhance the quality and/or nutritional value of fruits and vegetables.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts.

**Details of the AI usage are given below:**

1. ChatGPT

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Adekunle, A., Adeogun, O., and Olorunsuyi, Y. J. (2021). Effect of leaf extract of Lantana camara with maize-based coating on the quality of fresh-cut fruits of *Ananas comosus* and *Musa acuminata*. *Cogent Food & Agriculture*, 7(1), 1917834.
- Algarni, E. H. A., Elnaggar, I. A., El-wahed, N. A., Taha, I. M., AL-Jumay, H. A., Elhamamsy, S. M., Mahmoud, S. F., and Fahmy, A. (2022). Effect of chitosan nanoparticles as edible coating on the storability and quality of apricot fruits. *Polymers*, 14, 2227.
- Aloui, H., and Khwaldia, K. (2016). Natural antimicrobial edible coatings for microbial safety and food quality enhancement. *Comprehensive Reviews in Food Science and Food Safety*, 15(6), 1080–1103.
- Anean, H. A., Mallasiy, L. O., Bader, D. M., and Shaat, H.A. (2023). Nano edible coatings and films combined with zinc oxide and pomegranate peel active phenol compounds has been to extend the shelf life of minimally processed pomegranates. *Materials*, 16(4), 1569.
- APEDA [Agricultural and Processed Food Products Export Development Authority]. 2019. *Agriculture Export Policy*. Ministry of Commerce and Industry, GOI.
- Aytac, Z., Ipek, S., Durgun, E., Tekinay, T., and Uyar, T. (2017). Antibacterial electrospun zein nanofibrous web encapsulating thymol/cyclodextrin-inclusion complex for food packaging. *Food Chemistry*, 233, 117–124.
- Baek, J. H., Lee, S. Y., and Oh, S. W. (2021). Enhancing safety and quality of shrimp by nanoparticles of sodium alginate-based edible coating containing grapefruit seed extract. *International Journal of Biological Macromolecules*, 189, 84-90.
- Barrera-Necha, L. L., Correa-Pacheco, Z. N., Bautista-Banos, S., Hernández-Lopez, M., Jimenez, J. E. M., and Mejia, A. F. M. (2018). Synthesis and characterization of chitosan nanoparticles loaded botanical extracts with antifungal activity on *Colletotrichum gloeosporioides* and *Alternaria* species. *Advances in Microbiology*, 8, 286-296.
- Bautista-Banos, S., Barrera-Necha, L. L., Bravo-Luna, I., and Bermudes-Torres, L. (2002). Antifungal activity of leaf and stem extracts from various plant species on the incidence of *Colletotrichum gloeosporioides* of papaya and mango fruit after storage. *Revista Mexicana de Fitopatología*, 20, 8–12.
- Beltrán-Heredia, J., and Sánchez-Martín, J. (2009). Improvement of water treatment pilot plant with *Moringa oleifera* extract as flocculant agent. *Environmental Technology*, 30(6), 525–534.
- Bhutia, D. D., Zhimo, Y., Kole, R., and Saha, J. (2016). Antifungal activity of plant extracts against *Colletotrichum musae*, the postharvest anthracnose pathogen of banana cv. Martaman. *Nutrition & Food Science*, 46(1), 2–15.
- Cazón, P., Velazquez, G., Ramírez, J. A., and Vázquez, M. (2017). Polysaccharide-based films and coatings for food packaging: A review. *Food Hydrocolloids*, 68, 136–148.
- Chuang, P. H., Lee, C. W., Chou, J. Y., Murugan, M., Shieh, B. J., and Chen, H. M. (2007). Anti-fungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresource Technology*, 98(1), 232–236.
- Cordoba, P. L. J., and Sobral, P. J. A. (2017). Physical and antioxidant properties of films based on gelatin, gelatin-chitosan, or gelatin-sodium caseinate blends loaded with nanoemulsified active compounds. *Journal of Food Engineering*, 21, 347–353.
- Das, S. K., Vishakha, K., Das, S., Chakraborty, D., and Ganguli, A. (2022). Carboxymethyl cellulose and cardamom oil in a nanoemulsion edible coating inhibit the growth of foodborne pathogens and extend the shelf life of tomatoes. *Biocatalysis and Agricultural Biotechnology*, 42, 102369.
- Dave, R. K., Ramana Rao, T. V., and Nandane, A. S. (2017). Improvement of post-harvest quality of pear fruit with optimized composite edible coating formulations. *Journal of Food Science and Technology*, 54(12), 3917–3927.
- Deng, Z., Jung, J., Simonsen, J., and Zhao, Y. (2017). Cellulose nanomaterials emulsion coatings for controlling physiological activity, modifying surface morphology, and enhancing storability of postharvest bananas (*Musa acuminata*). *Food Chemistry*, 232, 359–368.
- Dhall, R. K. (2013). Advances in edible coatings for fresh fruits and vegetables: a review.



- Critical Reviews in Food Science and Nutrition*, 53(5), 435–450.
- Doughari, J. H., Elmahmoud, A. M., and Manzara, S. (2007). Studies on the antibacterial activity of root extracts of *Carica papaya* L. *African Journal of Microbiological Research*, 1, 37–41.
- Doughari, J. H., Pukuma, M. S., and De, N. (2007). Antibacterial effects of *Balanites aegyptiaca* L. Drel. and *Moringa oleifera* Lam. on *Salmonella Typhi*. *African Journal of Biotechnology*, 6(19), 2212–2215.
- El-Gioushy, S. F., Abdelkader, M. F., Mahmoud, M. H., Abou El Ghit, H. M., Fikry, M., Bahloul, A. M., Morsy, A. R., Abdelaziz, A. M., Alhaithloul, H. A., Hikal, D. M., and Abdein, M. A. (2022). The effects of a gum arabic-based edible coating on guava fruit characteristics during storage. *Coatings*, 12(1), 90.
- Emamifar, A., and Bavaisi, S. (2020). Nanocomposite coating based on sodium alginate and nano-ZnO for extending the storage life of fresh strawberries (*Fragaria xananassa* Duch.). *Journal of Food Measurement and Characterization*, 14, 1012–1024.
- Esyanti, R. R., Zaskia, H., Amalia, A., and Nugrahapraja, H. (2019). Chitosan nanoparticle-based coating as post-harvest technology in banana. *Journal of Physics: Conference Series*, 1204, 012109.
- Flores-López, M. L., Cerqueira, M. A., de Rodríguez, D. J., and Vicente, A. A. (2016). Perspectives on utilization of edible coatings and nano-laminate coatings for extension of postharvest storage of fruits and vegetables. *Food Engineering Reviews*, 8, 292–305.
- Gad, M., and Zog, Z. O. (2017). Mixing xanthan gum and chitosan nanoparticles to form a new coating to maintain storage life and quality of Elmamoura guava fruits. *International Journal of Current Microbiology and Applied Sciences*, 6(11), 1582–1591.
- Gurjar, M. S., Ali, S., Akhtar, M., and Singh, K. S. (2012). Efficacy of plant extracts in plant disease management. *Agricultural Sciences*, 3(3), 19046.
- Hager, J. V., Rawles, S. D., Xiong, Y. L., Newman, M. C., and Webster, C. D. (2019). Edible corn-zein-based coating incorporated with nisin or lemongrass essential oil inhibits *Listeria monocytogenes* on cultured hybrid striped bass, *Morone chrysops* x *Morone saxatilis*, fillets during refrigerated and frozen storage. *Journal of the World Aquaculture Society*, 50(1), 204–218.
- Hao, D. C., Gu, X. J., and Xiao, P. G. (2015). Medicinal plants: Chemistry, biology, and omics. *Woodhead Publishing*.
- Jianmei, L., Qi, S., Ying, S., Baorong, C., Xiaoling, W., and Tao, L. (2019). Improvement of banana postharvest quality using a novel soybean protein isolate/cinnamaldehyde/zinc oxide bionanocomposite coating strategy. *Scientia Horticulturae*, 258, 108786.
- Khaliq, G., Ramzan, M., and Baloch, A. H. (2019). Effect of Aloe vera gel coating enriched with *Fagonia indica* plant extract on physicochemical and antioxidant activity of sapodilla fruit during postharvest storage. *Journal of Food Chemistry*, 286, 346–353.
- Kubheka, S. F., Tesfay, S. Z., Mditshwa, A., and Magwaza, L. S. (2020). Evaluating the efficacy of edible coatings incorporated with moringa leaf extract on postharvest quality and its biofungicidal effect on 'Maluma' avocado fruit. *Horticultural Science*, 55(4), 410–415.
- Liu, C., Jin, T., Liu, W., Hao, W., Yan, L., and Zheng, L. (2021). Effects of hydroxyethyl cellulose and sodium alginate edible coating containing asparagus waste extract on postharvest quality of strawberry fruit. *LWT*, 148, 111770.
- Maringgal, B., Hashim, N., Tawakkal, I. S. M. A., Mohamed, M. T. M., Hamzah, M. H., and Ali, M. M. (2021). Effect of Kelulut honey nanoparticles coating on the changes of respiration rate, ascorbic acid, and total phenolic content of papaya (*Carica papaya* L.) during cold storage. *Foods*, 10, 432.
- Martelli, M., Barros, T. T., Moura, M. R., Mattoso, L. H. C., Assis, O. B. G. (2013). Effect of chitosan nanoparticles and pectin content on mechanical properties and water vapor permeability of banana puree films. *Journal of Food Science*, 78, 98–104.
- Melo, N. F. C. B., Lima, M. A. B., Stamford, T. L. M., Galembeck, A., Flores, M. A. P., Takaki, G. M. C., Medeiros, J. A. C., Stamford-Arnaud, T. M., and Stamford, T. C. M. (2020). In vivo and in vitro antifungal effect of fungal chitosan nanocomposite edible coating against strawberry phytopathogenic fungi. *International Journal of Food Science and Technology*, 55(11), 3381–3391.
- Mohammadi, A., Hashemi, M., and Hosseini, S. M. (2015). Chitosan nanoparticles loaded

- with *Cinnamomum zeylanicum* essential oil enhance the shelf life of cucumber during cold storage. *Postharvest Biology and Technology*, 110, 203–213.
- Ngubane, S., Tesfay, S. Z., Magwaza, L. S., and Mditshwa, A. (2024). The effect of composite edible coatings on the postharvest quality of “Hass” avocado fruit treated at different harvest maturities. *Frontiers in Sustainable Food Systems*, 8, 1473731.
- Nguyen, V. T. B., Nguyen, D. H. H., and Nguyen, H. V. H. (2020). Combination effects of calcium chloride and nano-chitosan on the postharvest quality of strawberry (*Fragaria x ananassa* Duch.). *Postharvest Biology and Technology*, 162, 111103.
- NHB [National Horticulture Board]. 2020. *Horticultural Statistics at a Glance*. Ministry of Agriculture & Farmers’ Welfare Department of Agriculture, Cooperation & Farmers’ Welfare, GOI.
- Nkurunziza, T., Nduwayezu, J. B., Banadda, E. N., and Nhapi, I. (2009). The effect of turbidity levels and *Moringa oleifera* concentration on the effectiveness of coagulation in water treatment. *Water Science and Technology*, 59(8), 1551–1558.
- Odetayo, T., Sithole, L., Shezi, S., Nomngongo, P., Tesfay, S., and Ngobesem, N. Z. (2022). Effect of nanoparticle-enriched coatings on the shelf life of Cavendish bananas. *Scientia Horticulturae*, 304, 111312.
- Otoni, C. G., Avena-Bustillos, R. J., Azeredo, H. M. C., Lorevice, M. V., Moura, M. R., and Mattoso, L. H. C. (2017). Recent advances on edible films based on fruits and vegetables—a review. *Comprehensive Reviews in Food Science and Food Safety*, 16(6), 1151–1169.
- Ozuna-Valencia, K. H., Moreno-Vásquez, M. J., Graciano-Verdugo, A. Z., Rodríguez-Félix, F., Robles-García, M. Á., Barreras-Urbina, C. G., Quintero-Reyes, I. E., Cornejo-Ramírez, Y. I., and Tapia-Hernández, J. A. (2024). The application of organic and inorganic nanoparticles incorporated in edible coatings and their effect on the physicochemical and microbiological properties of seafood. *Processes*, 12(9), 1889.
- Resende, N. S., Goncalves, G. A. S., Reis, K. C., Tonoli, G. H. D., and Boas, E. V. B. V. (2018). Chitosan/cellulose nanofibril nanocomposite and its effect on quality of coated strawberries. *Journal of Food Quality*, 13, 1727426.
- Saekow, M., Naradisorn, M., Tongdeesontorn, W., and Hamauzu, Y. (2019). Effect of carboxymethyl cellulose coating containing ZnO-nanoparticles for prolonging shelf life of persimmon and tomato fruit. *Journal of Food Science and Agricultural Technology*, 5, 41-48.
- Salvia-Trujillo, L., Rojas-Grau, M. A., Soliva-Fortuny, R., and Martin-Belloso, O. (2015). Use of antimicrobial nanoemulsions as edible coatings: Impact on safety and quality attributes of fresh-cut Fuji apples. *Postharvest Biology and Technology*, 105, 8–16.
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., and Latha, L. Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants’ extracts. *African Journal of Traditional, Complementary and Alternative Medicines*, 8(1), 1-10.
- Saucedo-Pompa, S., Torres-Castillo, J., Castro-Lopez, C., Rojas, R., Sanchez-Alejo, E., Ngangyo-Heya, M., and Martinez-Avila, G. (2018). Moringa plants: Bioactive compounds and promising applications in food products. *Food Research International*, 111, 438–450.
- Saxena, A., Sharma, L., and Maity, T. (2020). Enrichment of edible coatings and films with plant extracts or essential oils for the preservation of fruits and vegetables. In *Biopolymer-based formulations*, 859-880.
- Severino, R., Vu, K. D., Donsi, F., Salmieri, S., Ferrari, G., and Lacroix, M. (2014). Antibacterial and physical effects of modified chitosan-based coating containing nanoemulsion of mandarin essential oil and three non-thermal treatments against *Listeria innocua* in green beans. *International Journal of Food Microbiology*, 191, 82–88.
- Shafique, M., Rashid, M., Ullah, S., Rajwana, I. A., Naz, A., Razzaq, K., Hussain, M., Abdelgawad, M. A., El-Ghorab, A. H., Ghoneim, M. M., and Shaker, M. E. (2023). Quality and shelf life of strawberry fruit as affected by edible coating with moringa leaf extract, aloe vera gel, oxalic acid, and ascorbic acid. *International Journal of Food Properties*, 26(2), 2995–3012.
- Shankar, S., and Rhim, J. W. (2015). Amino acid mediated synthesis of silver nanoparticles and preparation of antimicrobial agar/silver

- nanoparticles composite films. *Carbohydrate Polymers*, 130, 353–363.
- Sharma, R., Nath, P.C., Das, P., Rustagi, S., Sharma, M., Sridhar, N., Hazarika, T.K., Rana, P., Nayak, P.K. and Sridhar, K. (2024). Essential oil-nanoemulsion based edible coating: Innovative sustainable preservation method for fresh/fresh-cut fruits and vegetables. *Food Chemistry*, 460, 140545.
- Soleimani, H., Mostowfizadeh-Ghalmfarsa, R., Ghanadian, M., Karami, A., and Cacciola, S.O. (2023). Defense mechanisms induced by celery seed essential oil against powdery mildew incited by *Podosphaera fusca* in cucumber. *Journal of Fungi*, 10(1), 17.
- Suryadi, Y., Samudra, M., and Soebrata, B. M. (2019). The use of curcumin-loaded chitosan nanoparticles to control anthracnose disease on papaya. *Suranaree Journal of Science and Technology*, 27(1), 1-12.
- Tabassum, P., Khan, S. A. K. U., Siddiqua, M., and Sultana, S. (2018). Effect of guava leaf and lemon extracts on postharvest quality and shelf life of banana cv. Sabri (*Musa sapientum* L.). *Journal of Bangladesh Agricultural University*, 16(3), 337–342.
- Tastan, O., Pataro, G., Donsi, F., Ferrari, G., and Baysal, T. (2017). Decontamination of fresh-cut cucumber slices by a combination of a modified chitosan coating containing carvacrol nanoemulsions and pulsed light. *International Journal of Food Microbiology*, 260, 75–80.
- Tesfay, S. Z., and Magwaza, L. S. (2017). Evaluating the efficacy of moringa leaf extract, chitosan, and carboxymethyl cellulose as edible coatings for enhancing quality and extending postharvest life of avocado (*Persea americana* Mill.) fruit. *Food Packaging and Shelf Life*, 11, 40–48.
- Unalan, I. U., Cerri, G., Marcuzzo, E., Cozzolino, C. A., and Farris, S. (2014). Nanocomposite films and coatings using inorganic nanobuilding blocks (NBB): Current applications and future opportunities in the food packaging sector. *RSC Advances*, 4, 29393.
- Vieira, A. C., Fonseca, J. M., Menezes, N. M. C., Monteiro, A. R., and Valencia, G. A. (2020). Active coatings based on hydroxypropyl methylcellulose and silver nanoparticles to extend the papaya (*Carica papaya* L.) shelf life. *International Journal of Biological Macromolecules*, 164, 489–498.
- Vongsak, B., Sithisarn, P., Mangmool, S., Thongpraditchote, S., Wongkrajang, Y., and Gritsanapan, W. (2013). Maximizing total phenolics, total flavonoids contents and antioxidant activity of *Moringa oleifera* leaf extract by the appropriate extraction method. *Industrial Crops and Products*, 44, 566–571.
- Yousuf, B., Wu, S., and Siddiqui, M. W. (2021). Incorporating essential oils or compounds derived thereof into edible coatings: Effect on quality and shelf life of fresh/fresh-cut produce. *Trends in Food Science & Technology*, 108, 245–257.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129675>